

On roots of Dehn twists.

Naoyuki Monden

Abstract

Margalit and Schleimer constructed nontrivial roots of the Dehn twist about a nonseparating curve. We prove that the conjugacy classes of roots of the Dehn twist about a nonseparating curve correspond to the conjugacy classes of periodic maps with certain conditions. Furthermore, we give data set which determine the conjugacy class of a root. As a consequence, we can find the minimum degree and the maximum degree, and show that the degree must be odd. Also, we give Dehn twist expression of the root of degree 3.

1 Introduction

Let $\Sigma_{g,b}$ denote a compact, oriented surface of genus g with b ($= 0, 2$) boundary components, and let $\mathcal{M}_{g,b}$ denote its mapping class group, the group of isotopy class of orientation-preserving homeomorphisms which are allowed to permute the points on the boundary. The isotopies are also required to permute the points on the boundary. If $b = 0$, we omit it from the notation. Let t_C be the (left handed) Dehn twist about a simple closed curve C on Σ_{g+1} ($g + 1 \geq 2$).

It is a natural question whether t_C has a root. In other words, does there exist $h \in \mathcal{M}_{g+1}$ such that for integer degree $n > 1$ $t_C = h^n$? If C is a separating curve, it is easy to find roots. For example, the half twist about C is a root of degree 2. However, when C is a nonseparating curve, roots were not discovered until recently. In 2009, Margalit and Schleimer [MS] discovered roots of the Dehn twist t_C about a nonseparating curve C of degree $2g + 1$.

Matsumoto and Montesinos [MM1] introduced a certain type of mapping class of Σ_{g+1} which were called by them pseudo-periodic maps of negative twist. Furthermore, they gave the complete set of conjugacy invariant for a pseudo-periodic map of negative twist. In Section 2 we introduce the definition and the complete set of conjugacy invariant of pseudo-periodic maps of negative twist. In Section 3 we show that inverses of roots of Dehn twists are pseudo-periodic maps of negative twist, and prove Main Theorem A by using the work of [MM1].

Main Theorem A. *Let C be a nonseparating curve. Let $\partial_\nu \Sigma_{g,2}$ ($\nu = 0, 1$) be the boundary of $\Sigma_{g,2}$, which we write $\overrightarrow{\partial_\nu \Sigma_{g,2}}$ to emphasize its orientation.*

The conjugacy classes ($\in \mathcal{M}_{g+1}$) of roots of t_C of degree n correspond to the conjugacy classes ($\in \mathcal{M}_{g,2}$) of periodic maps of order n which satisfy following condition:

The action on $\overrightarrow{\partial_\nu \Sigma_{g,2}}$ ($\nu = 0, 1$) is the rotation angle of $2\pi\delta^\nu/n$ such that $\delta^0 + \delta^1 = n - 1$ (Here $\gcd(\delta^\nu, n) = 1, 1 \leq \delta^\nu \leq n - 2$ ($\nu = 0, 1$)).

Moreover, by combining the Riemann-Hurwitz formula, Nielsen's theorem [Ni] and Main Theorem A we give the following data set which determines the conjugacy class of a root.

Main Theorem B. *The conjugacy class ($\in \mathcal{M}_{g+1}$) of a root h of t_C of degree n is determined by the data set $[n, g', (\sigma^0, \sigma^1); (\sigma_1, \lambda_1), \dots, (\sigma_k, \lambda_k)]$ which satisfies the following conditions:*

- (1) $2g/n = 2g' + \sum_{j=1}^k (1 - 1/\lambda_j)$,
- (2) $\sum_{j=1}^k \sigma_j n / \lambda_j + \sigma^0 + \sigma^1 \equiv 0 \pmod{n}$,
- (3) $\sigma^0 + \sigma^1 + \sigma^0 \sigma^1 \equiv 0 \pmod{n}$.

As a corollary of Theorem 3.3 we obtain the following result.

Corollary C. *the range of degree n is $3 \leq n \leq 2g + 1$ and n is odd.*

Thus the roots constructed by Margalit and Schleimer have the maximum degree $2g + 1$.

In Section 4 we give Dehn twist expression of the root of degree 3. In Section 5 we consider a root of image of the Dehn twist about a nonseparating curve to $\mathrm{Sp}(2(g + 1), \mathbb{Z})$. We show that the degree must be odd. In Section 6 we consider a root of the Dehn twist about a nonseparating curve in the mapping class group of a surface with boundary components and punctures.

Recently, McCullough and Rajeevsarathy [MR] has independently proved the result similar to Main Theorem A, B and Corollary C. They gave the data set similar to Theorem B in Theorem 1.1 and Main Theorem A was contained in the proof of Theorem 1.1. They proved these results without work of [MM1]. As a corollary of Theorem 1.1 they showed the same results as Corollary C. Moreover, they showed the follow results: In Corollary 3.1 they showed that t_C has a root of degree n for all $g + 1 > (n - 2)(n - 1)/2$. In Corollary 3.2 they showed that when n is prime and $g = (n - 2)(n - 1)/2$, t_c does not have a root of degree n (Corollary 3.2). In Theorem 4.2 they proved that if $n \geq g \geq 4$ then the root is one of 2 types of roots (Theorem 4.2).

The author thanks Darryl McCullough and Kashyap Rajeevsarathy for their comments on the content of this paper.

2 Preliminary

Hereafter, all surfaces will be oriented, and all homeomorphisms between them will be orientation-preserving. For us, a Dehn twist means a left-handed Dehn twist. Let Σ_{g+1} be a closed connected surface of genus $g + 1 \geq 2$.

2.1 Pseudo-periodic map and screw number

Definition 1. *A homeomorphism $f : \Sigma_{g+1} \rightarrow \Sigma_{g+1}$ is a pseudo-periodic map isotopic to a homeomorphism $f' : \Sigma_{g+1} \rightarrow \Sigma_{g+1}$ which satisfies the following conditions:*

- (1) *There exists a disjoint union of simple closed curves $\mathcal{C} = C_1 \cup C_2 \cup \dots \cup C_r$ on Σ_{g+1} such that $f'(\mathcal{C}) = \mathcal{C}$ ($\mathcal{C}$ might be empty).*

(2) The restriction $f'|_{(\Sigma_{g+1}-\mathcal{C})} : \Sigma_{g+1} - \mathcal{C} \rightarrow \Sigma_{g+1} - \mathcal{C}$ is isotopic to a periodic map.

We call $\{C_i\}_{i=1}^r$ an admissible system of cut curves if each connected component of $\Sigma_{g+1} - \mathcal{C}$ has negative Euler number. If Σ_{g+1} has negative Euler number, such a system always exists in the isotopy class of f .

Nielsen [Ni] introduced the "screw number" $s(C_i)$ for each component C_i of admissible system of cut curves $\mathcal{C}$. The definition is as follows:

We may assume $f(\mathcal{C}) = \mathcal{C}$. Let C_i be a fixed cut curve in the system.

Let α_i be the smallest positive integer α_i such that $f^{\alpha_i}(\vec{C}_i) = \vec{C}_i$ where $\vec{C}_i$ denotes C_i with an orientation assigned, so f^{α_i} preserves the (arbitrarily fixed) direction of C_i .

Let b_i and b'_i be the connected components of $\Sigma_{g+1} - \mathcal{C}$ such that C_i belongs to their adherence in Σ_{g+1} . Let β_i (resp. β'_i) be the smallest positive integer such that $f^{\beta_i}(b_i) = b_i$ (resp. $f^{\beta'_i}(b'_i) = b'_i$). Clearly α_i is a common multiple of β_i and β'_i .

Since $f|_{(\Sigma_{g+1}-\mathcal{C})}$ is isotopic to a periodic map, there exists a positive integer n_{b_i} such that $(f^{\beta_i}|_{b_i})^{n_{b_i}} \cong \text{id}|_{b_i}$. We choose the smallest number among such integers and denote it by n_{b_i} again. Likewise we choose $n_{b'_i}(> 0)$ for b'_i .

Let L_i be the least common multiple of $n_{b_i}\beta_i$ and $n_{b'_i}\beta'_i$. Then f^{L_i} is isotopic to the identity on b_i and b'_i . Thus, on the union $b_i \cup C_i \cup b'_i$, f^{L_i} is isotopic to the result of a number full Dehn twists performed about C_i . Let $e_i \in \mathbb{Z}$ be this number of full twists.

Definition 2. The rational number $e_i\alpha_i/L_i$ is called the screw number of f about C_i and is denoted by $s(C_i)$.

Definition 3. A pseudo-periodic map f is said to be of negative twist if $s(C_i) < 0$ for each component C_i of admissible system of cut curves $\mathcal{C}$.

Definition 4. The curve C_i is said to be amphidrome if α_i is even and $f^{\alpha_i/2}(\vec{C}_i) = -\vec{C}_i$, and non-amphidrome otherwise.

Theorem 5 ([MM1], Theorem 2.2). The conjugacy class of a negative twist $f : \Sigma_{g+1} \rightarrow \Sigma_{g+1}$ is determined by following data:

- An admissible system of cut curves $\mathcal{C} = \bigcup_i^r C_i$ on Σ_{g+1} .

For all $i = 1, \dots, r$,

- α_i : the smallest positive integer α_i such that $f^{\alpha_i}(\vec{C}_i) = \vec{C}_i$,
- the screw number $s(C_i)$ of f , and
- C_i 's character of being amphidrome or not.

For each connected component b of $\Sigma_{g+1} - \mathcal{C}$,

- β : the smallest positive integer such that $f^\beta(b) = b$,
- n_b : the smallest positive integer such that $(f^\beta|_b)^{n_b} \cong \text{id}|_b$, and
- the conjugacy class of a periodic map $f^\beta|_b$.
- The action of f on the oriented graph $G_{\mathcal{C}}$ whose vertices and edges correspond to connected components of $\Sigma_{g+1} - \mathcal{C}$ and $\{C_i\}_{i=1}^r$.

2.2 Valency

Let Σ be an oriented connected surface and let $f : \Sigma \rightarrow \Sigma$ be a periodic map of order $n > 1$. Let p be a point on Σ . There is a positive integer $\alpha(p)$ such that the points $p, f(p), \dots, f^{\alpha(p)-1}(p)$ are mutually distinct and $f^{\alpha(p)}(p) = p$. If $\alpha(p) = n$, we call the point p a simple point of f , while if $\alpha(p) < n$, we call p a multiple point of f . Note that a multiple point is an isolated and interior point of Σ .

Let $\vec{C} = \vec{C}'_1 \cup \vec{C}'_2 \cup \dots \cup \vec{C}'_s$ be a set of oriented and disjoint simple closed curves in a surface Σ , and let g be a map $g : \Sigma \rightarrow \Sigma$ such that $g(\vec{C}') = \vec{C}'$ and $g|_{\vec{C}'}$ is periodic. Let m_j be the smallest positive integer such that $g^{m_j}(\vec{C}'_j) = \vec{C}'_j$. The restriction $g^{m_j}|_{\vec{C}'_j}$ is a periodic map of $\vec{C}'_j$ of order, say, $\lambda_j > 0$. Then $m\lambda_j = n$ (= the order of $g|_{\vec{C}'}$). Let q be any point on $\vec{C}'_j$, and suppose that the images of q under the iteration of g^{m_j} are ordered as $(q, g^{m_j\sigma_j}(q), g^{2m_j\sigma_j}(q), \dots, g^{(\lambda_j-1)m_j\sigma_j}(q))$ viewed in the direction of $\vec{C}'_j$, where σ_j is an integer with $0 \leq \sigma_j \leq \lambda_j - 1$ and $\gcd(\sigma_j, \lambda_j) = 1$ ($j = 1, \dots, s$). So $\sigma_j = 0$ iff $\lambda_j = 1$. Let δ_j be the integer which satisfies

$$\sigma_j \delta_j \equiv 1 \pmod{\lambda_j}, \quad 0 \leq \delta_j \leq \lambda_j - 1 \quad (j = 1, \dots, s)$$

(NB. $\delta_j = 0$ iff $\lambda_j = 1$). Then the action of g^{m_j} on $\vec{C}'_j$ is the rotation of angle $2\pi\delta_j/\lambda_j$ with a suitable parametrization of $\vec{C}'_j$ as an oriented circle.

Definition 6 ([Ni]). *The triple $(m_i, \lambda_i, \sigma_i)$ and $(m_i, \lambda_i, \delta_i)$ are called the valency and the second valency of $\vec{C}'_j \in \vec{C}'$ with respect to g .*

Nielsen also defined the valency of a boundary curve as its valency with respect to f assuming it has the orientation induced by the surface Σ . The valency of a multiple point p is defined to be the valency of the boundary curve ∂D_p , oriented from the outside of a disk neighborhood D_p of p .

Let $\mathcal{C} = \bigcup_{i=1}^r C_i$ be an admissible system of cut curves subordinate to a pseudo-periodic map $f : \Sigma_{g+1} \rightarrow \Sigma_{g+1}$. We assume that $s(C_i)$ of f is non-zero ($i = 1, \dots, r$)

Thus Matsumoto and Montesinos proved the following proposition :

Proposition 7 ([MM1], Corollary 3.3.1, Corollary 3.7.1). *Let A_i be an annular neighborhood of C_i with $\partial A_i = \partial_0 A_i \cup \partial_1 A_i$. Let $(m_i^\nu, \lambda_i^\nu, \delta_i^\nu)$ be the second valency of $\partial_\nu A_i$ with respect to f ($\nu = 0, 1$). The second valency $(m_i^\nu, \lambda_i^\nu, \delta_i^\nu)$ of $\partial_\nu A_i$ oriented by the orientation induced from A_i is defined as the valency of $-\vec{C}_i$ with respect to the f .*

Then, If C_i is non-amphidrome,

- (1) $m_i^0 = m_i^1$,
- (2) $s(C_i) + \delta_i^0/\lambda_i^0 + \delta_i^1/\lambda_i^1$ is an integer.

If C_i is amphidrome,

- (1) $m_i^0 = m_i^1 =$ an even number,

- (2) $\delta_i^0 = \delta_i^1$ and $\lambda_i^0 = \lambda_i^1$,
(3) $s(C_i)/2 + \delta_i/\lambda_i$ is an integer,
(λ_i denotes $\lambda_i^0 = \lambda_i^1$, and δ_i denotes $\delta_i^0 = \delta_i^1$).

3 The conjugacy classes of roots of the Dehn twist about a nonseparating curve

3.1 Proof of Main Theorem A

In section 3.1 we do not distinguish a homeomorphism/curve and its isotopy class.

Let t_C be the Dehn twist about a simple closed curve C in Σ_{g+1} . Suppose that h is a root of t_C of degree $n > 1$. Since we have

$$t_C = h^n = hh^n h^{-1} = ht_C h^{-1} = t_{h(C)},$$

we see that $h(C) = C$. By $t_C = h^n$, $h|_{\Sigma_{g+1}-\{C\}}$ must be a periodic map of order n . Therefore h is a pseudo-periodic map, and an admissible system of cut curves is $\mathcal{C} = C$.

Let A be an annular neighborhood of C . Let $(m^0, \lambda^0, \delta^0)$ and $(m^1, \lambda^1, \delta^1)$ be the second valencies of $\partial_0 A$ and $\partial_1 A$ with respect to h .

Claim 8. *The simple closed curve C is non-amphidrome with respect to h .*

Proof. We assume that C is amphidrome with respect to h .

We will determine the screw number $s(C)$ of h . Let b be $\Sigma_{g+1} - C$. Let α , β and n_b be the smallest positive integers such that $h^\alpha(\vec{C}) = \vec{C}$, $h^\beta(b) = b$ and $(h^\beta|_b)^{n_b} = id_b$. Since $h(C) = C$ and C is amphidrome, we have $\alpha = m^0 = m^1 = 2$. Moreover, we have $\beta = 1$, $n_b = L$. Thus, by $h^n(\vec{C}) = t_C(\vec{C}) = \vec{C}$, we can denote $n = 2k$. By definition of $L > 0$, L is a divisor of n ($z := n/L \in \mathbb{Z}_{\geq 1}$). Then, we have $t_C = h^n = (h^L)^z = (t_C^e)^z = t_C^{ez}$. Therefore, we see that $e = 1$ and $L = 2k = n$. From the above arguments, we have

$$s(C) = 1/k.$$

By proposition 7, we have

$$\delta/\lambda = (2k - 1)/2k,$$

(Here λ denotes $\lambda^0 = \lambda^1$, and δ denotes $\delta^0 = \delta^1$.) However, since $n = 2k$ and the action of h^2 is the rotation of angle $2\pi\delta/\lambda$ in circle, δ/λ must be equal to δ/k . This is a contradiction. $\square$

Lemma 9. *Let C be a nonseparating simple closed curve in Σ_{g+1} . Then,*

$$\delta^0 + \delta^1 = n - 1 \quad (1 \leq \delta^\nu \leq n - 2, \nu = 0, 1)$$

Proof. We use Proposition 7. We will determine the screw number $s(C)$ of h . Since $h(C) = C$ and C is non-amphidrome (by Claim 8), we have $\alpha = m^0 = m^1 = 1$. Thus, we can find $s(C) = 1/n$ by the similar argument of Claim 8. Furthermore, we can find that $\lambda^\nu = \text{order of } h|_{\overrightarrow{\partial_\nu A}}$ (by $m^\nu \lambda^\nu = \text{order of } h|_{\overrightarrow{\partial_\nu A}}$ for $\nu = 0, 1$). By Proposition 7, we have $\delta^0/\lambda^0 + \delta^1/\lambda^1 = (n-1)/n$.

Let $\partial_\nu(\Sigma_{g+1} - A)$ be the boundary components which corresponds to $\partial_\nu A$ ($\nu = 0, 1$). Then, $h|_{(\Sigma_{g+1} - A)}$ is periodic map of order $n (= L)$. Therefore, we can get periodic map $\tilde{h} : \Sigma_g \rightarrow \Sigma_g$ of period n by pasting two disks D_0, D_1 to $\partial_0(\Sigma_{g+1} - A), \partial_1(\Sigma_{g+1} - A)$. Since A is non-amphidrome, we can see that $\tilde{h}$ fixes the center points D_0, D_1 . Consequently, the period λ^ν ($\nu = 0, 1$) of $h|_{\partial_\nu A}$ is equal to n . We note that δ^ν is not equal to 0. Because, if $\delta^\nu = 0$ then $n = \lambda^\nu$ is equal to 1 by the definition of λ^ν . This is in contradiction to $n > 1$. $\square$

Remark 10. If C is separating, then we may allow that λ^0 is not equal to λ^1 , that $\lambda^0, \lambda^1 < n$, and that either δ^0 or δ^1 is equal to 0. In other words, $(1, \lambda^0, \delta^0), (1, \lambda^1, \delta^1)$ satisfy any one of the following conditions :

- (1) $\delta^0/\lambda^0 + \delta^1/\lambda^1 = (n-1)/n$ ($\delta^0, \delta^1 \neq 0$) and $n = \text{lcm}(\lambda^0, \lambda^1)$,
- (2) $\delta^0 = 0, \delta^1 = n-1, \lambda^0 = 1$ and $\lambda^1 = n$,
- (3) $\delta^0 = n-1, \delta^1 = 0, \lambda^0 = n$ and $\lambda^1 = 1$.

Hereafter, C will be a nonseparating curve on Σ_{g+1} . Let $\Sigma_{g,2}$ be a compact oriented surface of genus g with two boundary components $\partial_0 \Sigma_{g,2}, \partial_1 \Sigma_{g,2}$.

Main Theorem A. The conjugacy classes $(\in \mathcal{M}_{g+1})$ of the roots of t_C of degree n correspond to the conjugacy classes $(\in \mathcal{M}_{g,2})$ of periodic maps of order n which satisfy following condition:

The action on $\overrightarrow{\partial_\nu \Sigma_{g,2}}$ ($\nu = 0, 1$) is the rotation angle of $2\pi\delta^\nu/n$ such that $\delta^0 + \delta^1 = n-1$ (Here $\gcd(\delta^\nu, n) = 1, 1 \leq \delta^\nu \leq n-2$ ($\nu = 0, 1$)).

Proof. We will prove Main Theorem A by using Theorem 5.

Let h be a root of Dehn twist t_C of degree n . Let G_C be the oriented graph G_C whose vertices and edges correspond to connected components of $\Sigma_{g+1} - C$ and C . Since C is non-amphidrome, we can find that the action of h on the oriented graph G_C is identity. From the above arguments, we have $\mathcal{C} = C$, $\alpha = 1, \beta = 1, n_b = n$, that C is non-amphidrome, and that the action of h on G_C is identity. These are the same data as h^{-1} .

Since the screw number $s(C)$ of h is equal to $1/n$, we see that $s(C)$ of h^{-1} is equal to $-1/n$. Therefore, h^{-1} is negative twist. If negative twists are restricted to roots h^{-1} of t_C^{-1} , the conjugacy class of a root h^{-1} of t_C^{-1} is determined by n and the conjugacy class of a periodic maps $h^{-1}|_{\Sigma_{g+1}-C}$ by using Theorem 5. It means that the conjugacy class of a root h of t_C is determined by n and the conjugacy class of a periodic map $h|_{\Sigma_{g+1}-C} = h|_{\Sigma_{g,2}}$. By Lemma 9 $h|_{\Sigma_{g+1}-C} = h|_{\Sigma_{g,2}}$ satisfies the condition.

In Section 3.2 we show that a root is constructed from a periodic map on $\Sigma_{g,2}$ which satisfies the condition. $\square$

3.2 Construction of a root of Dehn twist from a periodic map

Given a periodic map on $\Sigma_{g,2}$ with the condition of Main Theorem A, we construct a root of Dehn twist as follow:

Let $\tilde{f}$ be a periodic map on $\Sigma_{g,2}$ which satisfies the condition. Figure 1 shows the periodic map $\tilde{f}$.




Figure 1: The periodic map $\tilde{f}$

We will extend $\tilde{f}$ to Σ_{g+1} . Let us give an orientation to $[0, 1] \times S^1$ and $\partial_\nu \Sigma_{g,2}$ as the figure indicates:




Figure 2: $[0, 1] \times S^1$

Since $\tilde{f}|_{\partial_0 \Sigma_{g,2} \cup \partial_1 \Sigma_{g,2}}$ is periodic, there are homeomorphisms $\phi_0 : \{0\} \times S^1 \rightarrow \partial_0 \Sigma_{g,2}$ and $\phi_1 : \{1\} \times S^1 \rightarrow \partial_1 \Sigma_{g,2}$ such that $\tilde{f}\phi_0(0, x) = (0, x - \delta^0/n)$, $\tilde{f}\phi_1(1, x) = (1, x + \delta^1/n)$. Thus, we can get Σ_{g+1} by gluing $\{\nu\} \times S^1$ to $\partial_\nu \Sigma_{g,2}$ by ϕ_ν ($\nu = 0, 1$). Let ϕ_t be a homeomorphism $\phi_t : \{t\} \times S^1 \rightarrow \{t\} \times S^1$. We define f on Σ_{g+1} as follow:

$$\begin{aligned} f &= \tilde{f} \text{ on } \Sigma_{g,2} \\ f\phi_t(t, x) &= \phi_t(t, -(\delta^0/n)(1-t)x + (\delta^1/n)tx) \text{ on } [0, 1] \times S^1 \end{aligned}$$

Figure 3 shows a homeomorphism f .




Figure 3: the homeomorphism f

Figure 4 shows the homeomorphism $f|_{[0,1] \times S^1}$




Figure 4: $f|_{[0,1] \times S^1}$

Thus, we can see that $f^n \phi_t(t, x) = \phi_t(t, x + (n-1)t - \delta^0)$. Therefore, f^n is isotopic to t_C^{1-n} . So we may change f by isotopy so that $f^n = t_C^{1-n}$. By $f([0, 1] \times S^1) = [0, 1] \times S^1$, $ft_C = t_C f$. we have

$$t_C = (t_C f)^n$$

We have a root $(t_C f)$ of t_C with degree n . We complete the proof of Main Theorem A.

3.3 The data set which determines the conjugacy class of a root

Let p_j ($j = 1, \dots, k$) be a multiple point of a periodic map $\tilde{f} \in \mathcal{M}_{g,2}$ of Main Theorem A and let $(m_j, \sigma_j, \lambda_j)$ be the valency of p_j with respect to $\tilde{f}$. We note that $m_j \lambda_j = n$. Let $(1, \sigma^\nu, n)$ be the valency of $\partial_\nu \Sigma_{g,2}$ ($\nu = 0, 1$) respect to $\tilde{f}$. Note that $\sigma^\nu \delta^\nu \equiv 1 \pmod{n}$ and $1 \leq \sigma^\nu \leq n-1$ (by $n > 1$). By Nielsen's theorem [Ni], it suffices to classify the order of the map, the valencies of multiple points and the valencies of boundary curves.

When we combine the Riemann-Hurwitz formula, Nielsen's theorem and Main Theorem A, we have the following theorem:

Main Theorem B. *The conjugacy class $(\in \mathcal{M}_{g+1})$ of a root h of t_C of degree n is determined by the data $[n, g', (\sigma^0, \sigma^1); (\sigma_1, \lambda_1), \dots, (\sigma_k, \lambda_k)]$ which satisfies the following conditions:*

- (1) $2g/n = 2g' + \sum_{j=1}^k (1 - 1/\lambda_j)$,
- (2) $\sum_{j=1}^k \sigma_j n / \lambda_j + \sigma^0 + \sigma^1 \equiv 0 \pmod{n}$,
- (3) $\sigma^0 + \sigma^1 + \sigma^0 \sigma^1 \equiv 0 \pmod{n}$.

where g' is the genus of $\Sigma_{g,2}/\tilde{f}$.

McCullough and Rajeevsarathy also got the similar data set in [MR]. The notation $[n, g', (\sigma^0, \sigma^1); (\sigma_1, \lambda_1), \dots, (\sigma_k, \lambda_k)]$ follows that of [MR].

Corollary C. *the range of degree n is $3 \leq n \leq 2g+1$ and n is odd.*

Proof. If $n = 2$, by $1 \leq \sigma^0, \sigma^1 \leq n-1$ we have $\sigma^0 = \sigma^1 = 1$. This is in contradiction to the condition (3). Therefore, we see that n is odd.

For $n = 3$, if $g' = 0$, $k = g$ and $\sigma^0 = \sigma^1 = 1$, we can select σ_j ($j = 1, \dots, g$) which satisfy the condition (2). It means that there always exists the root of

degree 3 for $g \geq 1$. In next section, we will give Dehn twist expression of the root of degree 3.

We assume that $n > 2g + 1$. By the condition (1) we have $1 > 2g/n = 2g' + \sum_{j=1}^k (1 - 1/\lambda_j)$ so $g' = 0$ and $k = 1$. From the Condition (2) and (3) we have $n\sigma_1/\lambda_1 \equiv \sigma^0\sigma^1 \pmod{n}$. Therefore, we see that $0 \equiv n\sigma_1 \equiv \sigma^0\sigma^1\lambda_1 \pmod{n}$. It means that $\sigma^0\sigma^1\lambda_1/n = \sigma^0\sigma^1/m_1$ is an integer (we note that $m_1\lambda_1 = n$). Since $\gcd(\sigma^0, n) = \gcd(\sigma^1, n) = 1$ and $m_1\lambda_1 = n$, we see that m_1 must be equal to 1. It means that $\lambda_1 = n$. Thus we have $2g/n = 1 - 1/n$ so $n = 2g + 1$. This is in contradiction to $n > 2g + 1$. Since Margalit and Schleimer constructed the root of degree $2g + 1$, we have $n \leq 2g + 1$. $\square$

McCullough and Rajeevsarathy [MR] also proved by the similar arguments.

4 Dehn twist expression of the root of degree 3

We will give Dehn twist expression of the root of degree 3. The key points are to use the star relation which is given by Gervais [Ge].

Consider the torus with 3 boundary components d_1, d_2, d_3 and let a_1, a_2, a_3 and b be simple closed curves in Figure 5




Figure 5: The curves of star relation

The star relation is as follow:

$$(t_{a_1}t_{a_2}t_{a_3}b)^3 = t_{d_1}t_{d_2}t_{d_3}$$

If $a_1 = a_2$, then t_{d_3} is trivial, and the relation becomes

$$(t_{a_1}^2t_{a_3}b)^3 = t_{d_1}t_{d_2}$$

Let $\alpha_i, \alpha'_i, \beta_i$ ($i = 1, \dots, g+1$) and γ be nonseparating simple closed curves and let s_j ($j = 1, \dots, g-1$) be separating simple closed curves in Figure 6.




Figure 6: The curves $\alpha_i, \alpha', \beta_i$ ($i = 1, \dots, g+1$), γ, s_j ($j = 1, \dots, g-1$)

We define

$$\begin{aligned}\rho_1 &= (t_{\alpha_1} t_{\beta_1})^2 \\ \rho_i &= t_{\alpha_i}^2 t_{\alpha'_i} t_{\beta_i} \quad (i = 2, \dots, g-1) \\ \rho_g &= t_{\alpha_g} t_{\gamma} t_{\alpha'_g} t_{\beta_g}\end{aligned}$$

and

$$\begin{aligned}\hat{h} &= \rho_g \rho_{g-1}^{-1} \rho_{g-2} \cdots \rho_3^{-1} \rho_2 \rho_1^{-1} \quad (\text{if } g+1 \text{ is odd}), \\ \hat{h} &= \rho_g \rho_{g-1}^{-1} \rho_{g-2} \cdots \rho_3 \rho_2^{-1} \rho_1 \quad (\text{if } g+1 \text{ is even}).\end{aligned}$$

We note $\rho_1^3 = t_{s_1}$. Then, by star relation we have $\hat{h}^3 = t_{\alpha_{g+1}}^2$. When we define

$$h = t_{\alpha_{g+1}} \hat{h}^{-1},$$

h is the root of $t_{\alpha_{g+1}}$ of degree 3.

5 A root of an image of the Dehn twist about a nonseparating curve to $\mathrm{Sp}(2(g+1), \mathbb{Z})$

The action of $\mathcal{M}_{g+1}$ on $H_1(\Sigma_{g+1}; \mathbb{Z})$ preserves the algebraic intersection forms, so it induces a representation $\phi : \mathcal{M}_{g+1} \rightarrow \mathrm{Sp}(2(g+1), \mathbb{Z})$, which is well-known to be surjective. In this section we will prove that if C is nonseparating curve, then an image of Dehn twist t_C has no roots of even degree. We prove the case of $g+1 = 2$. An element 2×2 matrix $A \in \mathrm{Sp}(4, \mathbb{Z})$ satisfies that $AJ^t A = J$, where ${}^t A$ is transpose of A and J is

$$J = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}.$$

Let α_1 be a nonseparating simple closed curve in Figure 6 and let S be

$$S = \rho(t_{\alpha_1}) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (S^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}).$$

We assume that $S = A^2$, where $A \in \mathrm{Sp}(2(g+1), \mathbb{Z})$ is

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}.$$

Since $A^{-1} = S^{-1}A$, we have ${}^t A J = J A^{-1} = J S^{-1} A$. By

$${}^t A J = \begin{pmatrix} -a_{31} & -a_{41} & a_{11} & a_{21} \\ -a_{32} & -a_{42} & a_{12} & a_{22} \\ -a_{33} & -a_{43} & a_{13} & a_{23} \\ -a_{34} & -a_{44} & a_{14} & a_{24} \end{pmatrix}$$

and

$$JS^{-1}A = \begin{pmatrix} -a_{11} + a_{31} & -a_{12} + a_{32} & -a_{13} + a_{33} & -a_{14} + a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \\ -a_{11} & -a_{12} & -a_{13} & -a_{14} \\ -a_{21} & -a_{22} & -a_{23} & -a_{24} \end{pmatrix}$$

, we have

$$a_{11}/2 = a_{31}. \quad (1)$$

Since $SA = AS$, we have

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{11} + a_{31} & a_{12} + a_{32} & a_{13} + a_{33} & a_{14} + a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} = \begin{pmatrix} a_{11} + a_{13} & a_{12} & a_{13} & a_{14} \\ a_{21} + a_{23} & a_{22} & a_{23} & a_{24} \\ a_{31} + a_{33} & a_{32} & a_{33} & a_{34} \\ a_{41} + a_{43} & a_{42} & a_{43} & a_{44} \end{pmatrix}$$

We have $a_{12} = a_{13} = a_{14} = 0$. By $S = A^2$, we have

$$a_{11} = \pm 1. \quad (2)$$

By equations (1) and (2) we have that $a_{31} = a_{11}/2 = \pm 1/2$. This is in contradiction to $A \in \text{Sp}(4, \mathbb{Z})$. For $g > 2$, we can prove by the same arguments.

Since $S = \rho(t_C)$ has no roots of degree 2, we can find that a root of t_C cannot have even degree.

6 Remarks

We will consider a root of the Dehn twist about a nonseparating curve in the mapping class group of a surface with boundary components and punctures.

Let $\Sigma_{g,b_1,p}^{b_2}$ be a compact oriented surface with $b_1 + b_2$ boundary components $\partial_1, \dots, \partial_{b_1}$, $\partial'_1, \dots, \partial'_{b_2}$ and p punctures. We denote $\mathcal{M}_{g,b_1,p}^{b_2}$ the mapping class group of $\Sigma_{g,b_1,p}^{b_2}$, the group of isotopy classes of orientation-preserving homeomorphisms which are allowed to permute p punctures and the points on $\partial_1, \dots, \partial_{b_1}$, and restrict to the identity on $\partial'_1, \dots, \partial'_{b_2}$. The isotopies are also required to permute p punctures and the points on $\partial_1, \dots, \partial_{b_1}$ and to fix the points on $\partial'_1, \dots, \partial'_{b_2}$. Let t_C be the Dehn twist about a nonseparating curve C in $\Sigma_{g+1,b_1,p}^{b_2}$ ($g+1 \geq 2$). In this section we do not distinguish a homeomorphism/curve and its isotopy class.

We assume that $t_C = h^n \in \mathcal{M}_{g+1,b_1,p}^{b_2}$. Since $id|_{\Sigma_{g+1,b_1,p}^{b_2}-C} = t_C|_{\Sigma_{g+1,b_1,p}^{b_2}-C} = h^n|_{\Sigma_{g+1,b_1,p}^{b_2}-C}$, we see that $h|_{\Sigma_{g+1,b_1,p}^{b_2}-C}$ is a periodic map in $\mathcal{M}_{g,b_1+2,p}^{b_2}$. However, if $b_2 > 0$ then $\mathcal{M}_{g,b_1+2,p}^{b_2}$ is torsion free. This is in contradiction to what $h|_{\Sigma_{g+1,b_1,p}^{b_2}-C}$ is a periodic map in $\mathcal{M}_{g,b_1+2,p}^{b_2}$. Therefore, we see that if $b_2 > 0$, t_C has no roots in $\mathcal{M}_{g+1,b_1,p}^{b_2}$.

We assume that $b_2 = 0$ and omit b_2 from the notation.

Let $h(\in \mathcal{M}_{g+1})$ be a root of $t_C(\in \mathcal{M}_{g+1})$ of degree $2g+1$. By Main Theorem A and the proof of Corollary C, we see that $h|_{\Sigma_{g+1}-C} = h|_{\Sigma_{g,2,0}}$ is a periodic map of order $2g+1$ such that $h|_{\Sigma_{g,2,0}}$ maps $\partial_\nu \Sigma_{g,2,0}$ to $\partial_\nu \Sigma_{g,2,0}$ ($\nu = 0, 1$) and

has only one fixed point. Therefore, if $b_1 + p \equiv r \pmod{2g+1}$ and $1 < r < 2g+1$ then there is no $\mathbb{Z}_{2g+1}$ -action on $\Sigma_{g,b_1+2,p}$. It means that $t_C \in \mathcal{M}_{g+1,b_1,p}$ has no roots of degree $2g+1$. Therefore, we see that if $b_1 + p \equiv r \pmod{2g+1}$ and $1 < r < 2g+1$, $t_C \in \mathcal{M}_{g+1,b_1,p}$ has no roots of degree $2g+1$. In particular, if $g = 1$ and $b_1 + p \equiv 2 \pmod{3}$ then $t_C \in \mathcal{M}_{2,b_1,p}$ has no roots.

Reference

- [Ge] S. Gervais. *A finite presentation of the mapping class group of a punctured surface*. Topology, Vol. 40 (2001), No. 4, 703–725
- [MM1] Y. Matsumoto and J. M. Montesinos-Amilibia. *Pseudo-periodic maps and degeneration of Riemann surfaces*. I, II, preprint, Univ. of Tokyo and Univ. Complutense de Madrid, 1991/1992.
- [MM2] Y. Matsumoto and J. M. Montesinos-Amilibia. *Pseudo-periodic homeomorphisms and degeneration of Riemann surfaces*. Bull. Amer. Math. Soc. (N.S.) 30 (1994), no. 1, 70–75
- [MR] D. McCullough and K. Rajeevsarathy. *Roots of Dehn twists*. arXiv:0906.1601v1
- [Ni] J. Nielsen. *Surface Transformation Classes of Algebraically Finite Type*. Mat.-fys. Medd. Danske Vid.Selsk. 21 nr.2 (1944), [Collected Math. Papers, Vol.2, Birkhäuser, Boston, Basel, Stuttgart (1986)].

Department of Mathematics, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560-0043, Japan
E-mail address: **n-monden@cr.math.sci.osaka-u.ac.jp**